



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON THE ZEROS OF THE FUNCTION, $P(x)$, COMPLEMENTARY TO THE INCOMPLETE GAMMA FUNCTION*

BY

CHARLES N. HASKINS

INTRODUCTION

The problem of locating the real zeros of the function† $P(x)$, defined by the definite integral

$$P(x) \equiv \int_0^1 e^{-t} t^{x-1} dt, \quad x > 0,$$

or by the series

$$P(x) \equiv \sum_{s=1}^{s=\infty} \frac{(-1)^s}{s!} \frac{1}{x+s}$$

has been studied by Bourguet‡ whose results may be stated as follows:

The function $P(x)$ has no real zeros save in the intervals

$$\left. \begin{aligned} -2n < x < -2n + \frac{1}{2} \\ -2n + \frac{1}{2} < x < -2n + 1 \end{aligned} \right\}, \quad n = 3, 4, 5, \dots,$$

in each of which it has at least one.

The close relation of $P(x)$ to the gamma function, i. e., $\Gamma(x) = P(x) +$ integral transcendental function, and the fact that $P(x)$ is involved in certain functions used for the representation of statistical frequency distributions,§ make it desirable to complete Bourguet's results.

In this note I prove that the function $P(x)$ has at most two real zeros in each of the intervals

$$-2n < x < -2n + 1, \quad n = 3, 4, 5, \dots,$$

and consequently complete Bourguet's theorem so that it reads:

The function $P(x)$ has no real zeros save in the intervals

* Presented to the Society, January 2, 1915.

† Prym, F. E., *Journal für Mathematik*, vol. 82 (1876), pp. 165-172.

‡ Bourguet, L., *Acta Mathematica*, vol. 2 (1883), pp. 296-298.

§ Charlier, C. V. L., *Meddelande från Lunds Astronomiska Observatorium*, no. 26 (1905), p. 6.

$$-2n < x < -2n + \frac{1}{2}, \quad -2n + \frac{1}{2} < x < -2n + 1,$$

$$n = 3, 4, 5, \dots,$$

in each of which it has exactly one.

The proof depends upon an extension of the well-known theorem of Budan-Fourier. This extension though noticed by Stern* and Laguerre† and recently considered by Hurwitz,‡ appears not to have been employed in the literature to an extent commensurate with its merits.

1. The extended theorem of Budan-Fourier. In the interval $a \leq x \leq b$ let $f(x)$ have at each point a finite derivative of order N and let this derivative be of constant sign in the interval. Then the extended theorem of Budan-Fourier states that

The number of real roots of the equation $f(x) = 0$ which lie in the interval $a < x < b$ cannot be greater than the excess of the number of alternations of sign in the sequence

$$f(a), \quad f'(a), \quad f''(a), \quad \dots, \quad f^{(N-1)}(a), \quad f^{(N)}(a)$$

over that in the sequence

$$f(b), \quad f'(b), \quad f''(b), \quad \dots, \quad f^{(N-1)}(b), \quad f^{(N)}(b).$$

If the number of roots is not equal to this excess it falls short thereof by an even number.

The extended theorem may be proved by a method very similar to that used for the more restricted and usual form. The extended theorem is particularly useful in studying functions of the form $f(x) = \phi(x) + \phi_n(x)$, where $\phi_n(x)$ is a polynomial of degree n and $\phi(x)$ is a function whose N th derivative, $N > n$, is of constant sign in the interval under investigation.

2. Reduction to the interval $0 < x < 1$. The function $P(x)$ satisfies the difference-equation§

$$eP(x+1) = xeP(x) - 1.$$

If we consider the interval $-2n < x < -2n + 1$ and put $x = -2n + \theta$

$$P(x) = P(-2n + \theta), \quad 0 < \theta < 1.$$

Successive applications of the difference equation then give

$$eP(x) = eP(-2n + \theta) = \frac{\{eP(\theta) + W_{2n-1}(\theta)\}}{w_{2n}(\theta)},$$

* Stern, M. A., *Journal für Mathematik*, vol. 22.

† Laguerre, E., *Acta Mathematica*, vol. 4 (1884), p. 114.

‡ Hurwitz, A., *Mathematische Annalen*, vol. 71 (1911), pp. 584-591.

§ Prym, loc. cit., p. 167.

Now

$$W_3(\theta) = \theta^3 - 5\theta^2 + 9\theta - 4,$$

$$W_4(\theta) = \theta^4 - 9\theta^3 + 30\theta^2 - 41\theta + 20.$$

Hence we have

$$A_{k,k} = 1, \quad A_{k,r} > 3 > e, \quad k \geq 3, \quad r \leq k-1.$$

From the definition of $w_k(\theta)$ and $W_k(\theta)$ we obtain

$$w_k(\theta + 1) = \theta w_{k-1}(\theta),$$

$$W_k(\theta + 1) = 1 + \theta W_{k-1}(\theta),$$

from which we have

$$W'_k(\theta + 1) = W_{k-1}(\theta) + \theta W'_{k-1}(\theta),$$

$$W''_k(\theta + 1) = 2W'_{k-1}(\theta) + \theta W''_{k-1}(\theta),$$

$$\dots \dots \dots W^{(r)}_k(\theta + 1) = rW^{(r-1)}_{k-1}(\theta) + \theta W^{(r)}_{k-1}(\theta),$$

$$\dots \dots \dots W^{(k-1)}_k(\theta + 1) = (k-1)W^{(k-2)}_{k-1}(\theta) + \theta W^{(k-1)}_{k-1}(\theta),$$

$$W^{(k)}_k(\theta + 1) = kW^{(k-1)}_{k-1}(\theta);$$

and therefore, since

$$W^{(r)}_k(0) = (-1)^{k-r} A_{k,r} \cdot (r!),$$

$$W_k(1) = 1,$$

$$W'_k(1) = W_{k-1}(0) = (-1)^{k-1} A_{k-1,0},$$

$$W''_k(1) = 2W'_{k-1}(0) = (-1)^{k-2} A_{k-1,1} \cdot (2!),$$

$$\dots \dots \dots$$

$$W^{(r)}_k(1) = rW^{(r-1)}_{k-1}(0) = (-1)^{k-r} A_{k-1,r-1} \cdot (r!),$$

$$\dots \dots \dots$$

$$W^{(k-1)}_k(1) = (k-1)W^{(k-2)}_{k-1}(0) = -A_{k-1,k-2} \cdot \{(k-1)!\},$$

$$W^{(k)}_k(1) = kW^{(k-1)}_{k-1}(0) = k!.$$

From this we have at once the desired relations

$$W_k(1) = 1, \quad \text{sgn } W^{(r)}_k(1) = (-1)^{k-r}, \quad 1 \leq r \leq k; \quad W^{(k)}_k(1) = k!.$$

Since $A_{k-1,r} > 3 > e$, $k \geq 4$, $0 \leq r \leq k-2$, we have also

$$|W^{(r)}_k(1)| > e \cdot (r!), \quad 1 \leq r \leq k-1.$$

4. Certain properties of $P(\theta)$. Since

* Here $li(e^{-1})$ is the integral-logarithm, and C is Euler's constant, $0.57722 \dots$. Cf. Nielsen, *Theorie des Integrallogarithmus*, pp. 2, 11, 89.

are then evidently the same as those of the sequence (of $2n + 1$ terms)

$$P(\theta_0), P'(\theta_0), \dots P^{(r)}(\theta_0) \dots P^{(2n-1)}(\theta_0), P^{(2n)}(\theta_0),$$

and these are alternately positive and negative. There are therefore $2n$ alternations of sign in this sequence.

On passing to the other end of the interval we have

$$R_{2n}(1) = eP(1) + W_{2n-1}(1) = +e,$$

$$R'_{2n}(1) = eP'(1) + W'_{2n-1}(1) = +\{A_{2n-2,0} - 2.16538 \dots\},$$

$$R''_{2n}(1) = eP''(1) + W''_{2n-1}(1) = -\left\{A_{2n-2,1} - \frac{e|P''(1)|}{2!}\right\}(2!),$$

.

$$R^{(r)}_{2n}(1) = eP^{(r)}(1) + W^{(r)}_{2n-1}(1) = (-1)^{r-1}\left\{A_{2n-2,r-1} - \frac{e|P^{(r)}(1)|}{r!}\right\}(r!),$$

.

$$\begin{aligned} R^{(2n-2)}_{2n}(1) &= eP^{(2n-2)}(1) + W^{(2n-2)}_{2n-1}(1) \\ &= -\left\{A_{2n-2,2n-3} - \frac{e|P^{(2n-2)}(1)|}{(2n-2)!}\right\}(2n-2)!, \end{aligned}$$

$$R^{(2n-1)}_{2n}(1) = eP^{(2n-1)}(1) + W^{(2n-1)}_{2n-1}(1) = -\left\{\frac{e|P^{(2n-1)}(1)|}{(2n-1)!} - 1\right\}(2n-1)!,$$

$$R^{(2n)}_{2n}(1) = eP^{(2n)}(1) = +|eP^{(2n)}_{(1)}|.$$

Now if $n \geq 3$, $2n - 2 \geq 4$. Hence

$$A_{2n-2,0} > 3 > 2.16538 \dots, \quad A_{2n-2,r-1} > e > \frac{e|P^{(r)}(1)|}{r!}, \quad (1 \leq r \leq 2n-2),$$

but

$$\frac{e|P^{(2n-1)}(1)|}{(2n-1)!} - 1 > 0.$$

Hence the sequence of signs in

$$R_{2n}(1), R'_{2n}(1), R''_{2n}(1), \dots R^{(r)}_{2n}(1), \dots R^{(2n-3)}_{2n}(1), R^{(2n-2)}_{2n}(1),$$

$$R^{(2n-1)}_{2n}(1), R^{(2n)}_{2n}(1)$$

is

$$+ \quad + \quad - \quad \dots \quad (-1)^{r-1} \dots \quad + \quad - \quad - \quad +$$

As compared with the sequence of signs at $\theta = \theta_0$ we see that this sequence has lost two alternations and only two, namely, one between R and R' , and one between $R^{(2n-2)}$ and $R^{(2n-1)}$.

Hence $R_{2n}(\theta)$ can have at most two zeros in the interval $\theta_0 \leq \theta \leq 1$ and therefore at most two in the interval $0 < \theta < 1$.

Therefore $P(x)$ can have at most two zeros in the interior of the interval

$$-2n < x < -2n + 1;$$

which is the theorem to be proved.

DARTMOUTH COLLEGE,
December 14, 1914.
